



PAEDIATRIC ACUTE CARE GUIDELINE

Hypothermia

Scope (Staff):	All Emergency Department Clinicians
Scope (Area):	Emergency Department

This document should be read in conjunction with this **DISCLAIMER**
<http://kidshealthwa.com/about/disclaimer/>

Hypothermia

Hypothermia is core temperature < 35°C. Young children are at risk due to high body surface area to weight ratio.

Background

- Hypothermia usually due to environmental causes e.g. immersion and exposure
- Check core temperature using a rectal or oesophageal thermometer
- Be aware that sepsis may present with hypothermia

General

Effects of hypothermia

32°C	<ul style="list-style-type: none"> • Decreased pulse / respiratory rate / BP / conscious state • Shivering stops. Muscle rigidity (may mimic rigor mortis) • Atrial arrhythmias appear - usually innocent and revert when rewarmed
28°C	<ul style="list-style-type: none"> • Ventricular arrhythmias (including VF) • Fixed dilated pupils
26°C	<ul style="list-style-type: none"> • Comatose • Absent reflexes • Apnoea
20°C	<ul style="list-style-type: none"> • Asystole
15°C	<ul style="list-style-type: none"> • Temperature of the coldest known survivor

Hypothermia in Resuscitation

- Hypothermia substantially reduces effectiveness of defibrillation and resuscitation drugs. It is reasonable to attempt defibrillation, but if unsuccessful, continue cardiac compression until core temperature is $> 30^{\circ}\text{C}$, when defibrillation / drugs are more likely to be effective.
- Drugs are generally withheld until core temperature is $>30^{\circ}\text{C}$, as accumulation may occur while cold, with resultant toxicity when rewarmed.
- Never diagnose death and thus stop resuscitation until the patient is rewarmed to at least 32°C or cannot be rewarmed despite active measures.

Assessment

- Monitor core temperature
- Monitor heart rate and rhythm
- Check electrolytes and glucose

Initial management

Important Principles:

- Actively rewarm to 32°C , then allow passive rewarming. Once above the fibrillation threshold (32°C) there is no urgency in rewarming
- Mild brain hypothermia may limit reperfusion injury
- Avoid hyperthermia (keep temperature $< 36.5^{\circ}\text{C}$)
- Never diagnose death and thus stop resuscitation until the patient is rewarmed to at least 32°C , or cannot be rewarmed despite active measures
- Beware: rewarming may lead to vasodilation and hypotension (so-called "*after shock*"), which can contribute significantly to mortality
- Beware: Peripheral rewarming and vasodilation can result in cold, acidotic blood being shunted to the core, with a drop in core temperature (so-called "*after drop*") and an increased risk of arrhythmias
- Hypokalaemia is common, even in the presence of marked acidosis
- Check blood gases, potassium, glucose, and haematocrit with every few degrees of warming

Further management

External rewarming (for temperature $> 32^{\circ}\text{C}$)

- Passive external rewarming:
 - Remove wet clothes, dry patient
 - Warm blankets
 - Cover with sheet of foil/space blankets

- Active external rewarming (truncal areas only):
 - Overhead warmers
 - Warm air system e.g. Bair Hugger
 - Thermal mattresses


Active core rewarming (for temperature < 32°C)

- Warm IV fluids to 39°C with blood warmer (slow) – start with pre-warmed IV 0.9% saline at 40°C
- Gastric or bladder lavage with 0.9% saline at 40°C
- Peritoneal lavage with potassium-free dialysate or 0.9% saline at 40°C. Use 20 mL/kg cycled every 15 minutes
- Ventilation with humidified gas heated to 42°C
- Pleural or pericardial lavage
- Haemodialysis, extra-corporeal blood warming

Nursing

Routine nursing care.

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Dr Meredith Borland HoD, PMH Emergency Department		
Reviewer / Team:	Kids Health WA Guidelines Team		
Date First Issued:	12 November, 2014	Version:	
Last Reviewed:	5 November, 2015	Review Date:	5 November, 2017
Approved by:	Dr Meredith Borland	Date:	5 November, 2015
Endorsed by:	Medical Advisory Committee	Date:	5 November, 2015
Standards Applicable:	NSQHS Standards: 		
Printed or personally saved electronic copies of this document are considered uncontrolled			