



PAEDIATRIC ACUTE CARE GUIDELINE

Peritonsillar Abscess

Scope (Staff):	All Emergency Department Clinicians
Scope (Area):	Emergency Department

This document should be read in conjunction with this DISCLAIMER
<http://kidshealthwa.com/about/disclaimer/>

Peritonsillar Abscess

A peritonsillar abscess (also called paratonsillar abscess) is a collection of pus in the space between the tonsil and the superior pharyngeal constrictor muscle.

Background

A peritonsillar abscess is often considered to be a complication of tonsillitis. However, it is now thought to be secondary to infection of a peritonsillar salivary gland (Weber gland) which is located between the tonsillar capsule and the muscle of the tonsillar fossa.

Assessment

Patients often present with:

- Severe sore throat
- Odynophagia (painful swallowing) with drooling
- Muffled voice ('Hot Potato Voice')
- Difficulty opening mouth (trismus)

Examination often reveals:

- Limited mouth opening (trismus), less than three finger width
- Unilateral swollen enlarged tonsil with fluctuant swelling extending up to the soft palate (most characteristic)
- Deviation of the uvula away from the affected side

- Enlarged tender cervical lymph node on the associated side
- The patient is usually febrile, and often 'toxic' looking

Management

- Patients require hospitalisation for rehydration, intravenous antibiotics, analgesia and, in most cases, surgical drainage of the abscess.
- Antibiotics need to cover *Streptococcus pyogenes* and anaerobes
 - A combination of intravenous Benzylpenicillin and Metronidazole is recommended - [Antibiotics](#)
- As a rule of thumb, all peritonsillar abscesses should be drained
 - Generally, children less than 7-10 years of age will not tolerate oropharyngeal procedures under local anaesthetic very well. Needle aspiration or incision and drainage of the abscess under general anaesthesia is usually required.
 - Patients who are septic and have airway obstruction may be considered for quinsy tonsillectomy.

It is worth noting that in some very young children quinsy tends to resolve with IV antibiotics, hence medical treatment and observation for 24 hours may be worthwhile.

Nursing

- Baseline observations: heart rate, respiratory rate, temperature, SpO2, blood pressure and pain score
- Minimum of hourly observations should be recorded whilst in the Emergency Department
- Fluid input/output is to be monitored and documented


Tags

abscess, airway, benzylpenicillin, cephazolin, clindamycin, dehydration, deviation, drooling, enlarged, febrile, fever, flagyl, glan, glans, lymph, metronidazole, mouth, neck, node, nodes, odynophagia, oedema, pain, painful, parapharyngeal, paratonsillar, paratonsillar abscess, penicillin, peritonsillar, peritonsillar abscess, pharyngeal, pus, quinsy, sepsis, sore throat, swelling, swollen, tonsil, tonsillar, tonsillitis, tonsils, trismus, uvula, weber gland

References

1. Wald ER (2014) Peritonsillar Cellulitis and Abscess. UpToDate. Accessed at www.uptodate.com
2. WA Health Child and Adolescent Health Service. Ear, Nose, Throat and Dental ChAMP Empiric Guidelines Version 2, February 2014

This document can be made available in alternative formats on request for a person with a disability.

File Path:			
Document Owner:	Dr Meredith Borland HoD, PMH Emergency Department		
Reviewer / Team:	Kids Health WA Guidelines Team		
Date First Issued:	7 May, 2015	Version:	
Last Reviewed:	7 May, 2015	Review Date:	7 May, 2017
Approved by:	Dr Meredith Borland	Date:	7 May, 2015
Endorsed by:	Medical Advisory Committee	Date:	7 May, 2015
Standards Applicable:	NSQHS Standards: 		
Printed or personally saved electronic copies of this document are considered uncontrolled			