# Princess Margaret Hospital for Children Emergency Department Guideline

PAEDIATRIC ACUTE CARE GUIDELINE			
Kawasaki Disease			
Scope (Staff):	All Emergency Department Clinicians		
Scope (Area):	Emergency Department		

This document should be read in conjunction with this DISCLAIMER <a href="http://kidshealthwa.com/about/disclaimer/">http://kidshealthwa.com/about/disclaimer/</a>

## Kawasaki Disease

## **Background**

- Kawasaki disease is a common vasculitis of childhood especially in < 5 year olds
- Aetiology is unknown
- Is a self limiting condition, with fever and manifestations of acute inflammation lasting an average 12 days without therapy, however, early treatment is necessary to prevent complications

## **Complications**

- Coronary artery aneurysms
  - Infants under 12 months at increased risk of coronary artery aneurysm
  - Delay of treatment (after 10 days) increases risk of coronary artery aneurysm by five times
- · Depressed myocardial contractility & heart failure
- Myocardial infarction
- Arrhythmias
- Peripheral arterial occlusion

## **Assessment**

## **Diagnostic Criteria**

Kawasaki Disease is a clinical diagnosis with no diagnostic laboratory test.

Presence of prolonged unexplained fever ≥ 5 days (fever > 38.5°C) with at least 4 of the

#### following criteria:

- 1. Bilateral non-exudative conjunctivitis
- 2. Polymorphous rash
- 3. Cervical lymphadenopathy (at least 1 lymph node >1.5cm in diameter)
- 4. Mucositis cracked red lips, injected pharynx or strawberry tongue
- 5. Extremity changes erythema of palms/soles, oedema of hands/feet (acute phase), and periungual desquamation (convalescent phase)

#### Associated non-specific symptoms

- Diarrhoea, vomiting, or abdominal pain
- Irritability
- Cough or rhinorrhea
- Joint pain
- Weakness

#### **Incomplete (Atypical) Kawasaki Disease**

- Diagnostic criteria not completely fulfilled (< 4 signs of mucocutaneous inflammation)</li>
   but otherwise similar clinical picture to that of "classic" Kawasaki Disease
- More likely with children < 12 months old and > 5 years old
- Atypical Kawasaki Disease patients are still at risk of cardiovascular complications
  - If prolonged unexplained fever discuss with ED Consultant/on call General Paediatric Consultant

## Laboratory findings (not diagnostic but supportive)

- Elevated acute phase reactants (CRP, ESR)
- Elevated WCC with predominant neutrophilia
- Elevated platelets (after 1 week)
- Normocyctic, normochromic anaemia
- Sterile pyuria (need clean voided specimen)

### Investigations

No diagnostic lab tests for Kawasaki Disease but can be supportive or used to exclude other causes of fever.

- CPR, ESR, FBC, ALT, Albumin
- ASOT/AntiDNAse B
- Urinalysis clean catch or in-out catheter
- Blood culture

#### **Differential diagnoses**

- Measles
- Adenovirus
- Epstein Barr Virus
- Scarlet fever
- Toxic Shock Syndrome
- Steven-Johnson Syndrome

## **Management**

All suspected cases should be discussed with the ED Admitting Registrar/Consultant for admission under the on call General Paediatric Consultant

#### **Initial Management**

- IV Immunoglobulin (IVIG) 2 grams/kg over 8-12 hours
- Low dose aspirin at 3-5mg/kg daily

#### **Further Management**

- Second dose IVIG may be given if incomplete treatment response, particularly in high risk age group and "atypical" Kawasaki Disease
- General Paediatric Team will refer the patient to Cardiology for echocardiogram **only after** the diagnosis of Kawasaki Disease is made/confirmed and treatment instituted
- Many patients (especially if < 3 years old) will require sedation to perform the echocardiogram as the irritability (commonly seen) precludes performing adequate echocardiogram in the acute phase
- Echocardiogram is required at/after initial diagnosis and repeated at 4-8 weeks post treatment
- Echocardiogram plays no role in the diagnosis of Kawasaki Disease
- Referral to the Infectious Disease Consultant is at the discretion of the General Paediatric Consultant

## **Discharge Treatment**

 Continue aspirin (low dose) at 3-5mg/kg daily until the repeat echocardiogram at 4-8 weeks confirms absence of coronary involvement

# Nursing

Routine nursing care.

#### References

- 1. Sundal R (2014) Kawasaki Disease: Clinical Features and Diagnosis. UpToDate. Accessed at www.uptodate.com
- 2. Sundal R (2014) Kawasaki Disease: Initial Treatment and Prognosis. UpToDate. Accessed at www.uptodate.com
- 3. AMH Children's Dosing Companion (online). Adelaide: Australian Medicines Handbook Pty Ltd; 2015 January. Available from:

https://childrens-amh-net-au.pklibresources.health.wa.gov.au

This document can be made available in alternative formats on request for a person with a disability.

File Path:				
Document Owner:	Dr Meredith Borland HoD, PMH Emergency Department			
Reviewer / Team:	Kids Health WA Guidelines Team			
Date First Issued:	2 June, 2015	Version:		
Last Reviewed:	2 June, 2015	Review Date:	2 June, 2017	
Approved by:	Dr Meredith Borland	Date:	2 June, 2015	
Endorsed by:	Medical Advisory Committee	Date:	2 June, 2015	
Standards Applicable: NSQHS Standards:				

Printed or personally saved electronic copies of this document are considered uncontrolled